

Condition Treated	Active Health Avg.% Improvement	ChiroUp Avg.% Improvement	Active Health Avg. # of Visits	ChiroUp Avg. # of Visits
<b>Overall</b>	<b>64.08%</b>	<b>71.1%</b>	<b>5.6</b>	<b>6.8</b>
Acute Cervical Torticollis	—	80.4%	—	4.9
Biceps Tendinopathy- Proximal	80%	68.1%	3	7.6
Cervical Disc Lesion	—	66.7%	—	8.6
Cervical Segmental Joint Restriction	—	69.5%	—	6.7
Cervicogenic Headache	—	72.6%	—	6.9
Chondromalacia Patellae	—	62.6%	—	5.9
Costovertebral Dysfunction	—	80.2%	—	5.6
Dysfunctional Breathing	—	71.9%	—	6
Foot Hyperpronation	26.7%	55%	2	5.5
Greater Trochanteric Pain Syndrome	—	75.8%	—	7.2
Hip Abductor Weakness	—	67.8%	—	6.4
Iliotibial Band Syndrome	100%	74.4%	7	7.1
Knee Sprain- Collateral Ligament	100%	77.1%	7	7.1
Lateral Epicondylopathy	50%	69.2%	8	6.5
Lower Crossed Syndrome	70%	70.4%	7	6.3
Lumbar Disc Lesion	66.7%	71.5%	5.7	8.2
Lumbar Facet Syndrome	92.5%	70.8%	9.2	7
Lumbar Segmental Joint Restriction	—	72.9%	—	6.8
Lumbar Spondylolisthesis- Type II Isthmic	—	65%	—	6.8
Lumbar Spondylolysis	—	74%	—	6.8
Lumbar Spondylosis/ DJD/ DDD	—	72.1%	—	7.7
Lumbar Sprain/ Strain	95%	81.2%	3	6.8
Medial Tibial Stress Syndrome	26.7%	59.2%	2	5.8
Modifier- Cervical Radiculopathy	—	61%	—	7
Modifier- Extension Biased LBP	64%	73.6%	5.6	6.8
Modifier- Lumbar Radiculopathy	—	65.7%	—	8.4
Plantar Fasciitis	—	54.7%	—	5.8
Rotator Cuff Strain/ Tendinopathy	70%	62.5%	9	6.4
Scapular Dyskinesis	80%	70%	3	7
Scoliosis	—	58.8%	—	5.7
Shoulder Anterior Impingement Syndrome	—	68.4%	—	6
Thoracic Outlet Syndrome	—	70.9%	—	7.4
Upper Crossed Syndrome	30%	70.6%	7	6.7